## **Sublingual Immunotherapy**



Patient:	DOB:	Doctor:

## **MAINTENANCE SCHEDULE**

<u>Directions</u>: Each dose should be applied under the tongue and held there for two minutes. Swallow the dose afterwards. \*\*If local mouth itching occurs, then spit out the dose.\*\* Please report any symptoms under "Remarks", such as itching in mouth, swelling under tongue, upset stomach, rashes, etc.

Each vial contains a 1.5 month- supply. Take 0.25 mL (5 drops) once daily.

POLLEN VIAL			DME VIAL		
Date	Remarks	Date	Remarks		

POLLEN VIAL		DME VIAL			
Date	Remarks	Date	Remarks		
	_				
You now have a	You now have approximately 2 weeks of drops left. Please call to reorder next vial(s).				

## Things to know:

- Refrigerate your vials. Call our office to reorder within two weeks before they are empty.
- Always take the same dose from each vial. We want the vials to empty at the same time.
- **Missed doses:** 1 week take the next dose when you remember. 2 + weeks contact us.
- All patients are required to see the doctor twice a year. Your last office visit was\_\_\_\_\_
- Should you have a reaction to your allergy drops (tickle in your throat, itching all over, ect):
  - Take a Benadryl (Adult 50 mg or 4 tsp; Child 25 mg or 2 tsp) or Liquid Zyrtec followed by Zantac (Adult 150 mg; Child 75 mg)
- Should symptoms worsen (throat closing up, tightness in chest, etc.):
  - Use your Epi-Pen and Dial 911